

DENTAL IMPLANTS INSTRUCTIONS

The below instructions will cover proper post-surgical care for dental implants.

SINUS PRECAUTIONS:

General Reminders:

- Do not disturb the surgical site.
- Avoid touching the area or spitting for a few days following surgery.
- There may be a metal spacer slightly protruding through the gum tissue.

BLEEDING:

- A small amount of bleeding is common for the first few days.
- If you experience excessive bleeding, place a gauze pad directly over the surgical sites and hold it in place with firm biting pressure for 30–45 minutes or until the bleeding can be controlled.
- Replace the gauze pads every 30–45 minutes.
- Bleeding may vary from person to person.
- Gauze pads will always appear red. The way to assess bleeding is to directly look at the sites after the gauze pads have been removed.
- If the bleeding does not slow down, please call our office.

SWELLING CONTROL:

- Swelling can be normal after surgery.
- Swelling can be reduced by applying an ice pack to the side of your face. Do this for 20 minutes on, 20 minutes off.
- Continue icing the face as much as possible for the first 24–36 hours. Do not apply ice directly to the skin.
- Please do not lay flat. Use 1–2 pillows to elevate the head to avoid or minimize swelling.
- These measures may not eliminate swelling but they will help reduce its severity.

DIET-GENERAL:

- Drink plenty of fluids.
- Avoid hot liquids or hot foods. Please refer to the post-op diet instructions provided by our office.

DIET-DAY OF PROCEDURE:

- Soft liquidy foods are strongly recommended on the day of the procedure. Pudding, JELL-O®, applesauce, yogurt, ice cream, and smoothies are excellent options for the day of the procedure.

DIET-FIRST WEEK:

- You should eat only soft foods for the first week, such as scrambled eggs, mashed potatoes, macaroni and cheese, soft fish, and steamed vegetables.
- Advance your diet as tolerated after the initial week.

DIET-FOODS TO AVOID:

- Avoid hard, crunchy, or very chewy foods for the first 8 weeks following surgery.

PAIN CONTROL:

- Discomfort can be normal after the procedure.
- Pain medication may be prescribed. It is advised that you take pain medication on the first day.
- If you are not allergic or intolerant to nonsteroidal anti-inflammatory drugs, you may start taking ibuprofen (Advil® or Motrin®).
- If the ibuprofen alone doesn't control your discomfort, take the prescribed narcotic in addition.
- It is recommended that you alternate between the narcotic and ibuprofen. Do not take them at the same time.
- Be certain to take your pain medication with food, as this will help prevent an upset stomach.
- Remember: narcotic pain medication will impair your judgment and reflexes, so do not drive or operate heavy machinery.

ALCOHOLIC BEVERAGES & SMOKING:

- Alcoholic beverages should be avoided for 24 hours before and 5 days after surgery.
- Smoking should be avoided or limited as much as possible.



DENTAL IMPLANTS INSTRUCTIONS (CONT.)

MOUTH RINSES:

- If a mouth rinse has been prescribed, please use it as directed.
- Warm salt water rinses can be made by mixing ½ teaspoon of salt with a cup of warm water and should be used at least 4-5 times a day, especially after meals.

HYGIENE:

- You are advised to brush your teeth normally; however, please avoid the surgical site the day of the procedure.
- Be gentle initially with brushing the surgical areas. Gently brushing the healing abutment is encouraged.

PHYSICAL ACTIVITY:

- No strenuous activity for 3-4 days following your surgery.
- Physical activity increases your blood pressure, which will cause an increase in your swelling, pain, and bleeding.
- You may gradually increase your activity after 3-4 days.
- Low impact activities, such as swimming, may be more comfortable at first.
- Avoid bending over, heavy lifting, or straining.

Thank you for reviewing these post-operative instructions and following them carefully.